

Abdominal Trauma

WWW.RN.ORG®

Reviewed December, 2020, Expires December, 2022
Provider Information and Specifics available on our Website
Unauthorized Distribution Prohibited

©2020 RN.ORG®, S.A., RN.ORG®, LLC


By Wanda Lockwood, RN, BA, MA

Purpose The purpose of this course is to outline the different types of abdominal injuries that may result from blunt or penetrating trauma, including diagnosis, symptoms, associated injuries, and treatment approaches.

Goals Upon completion of this course, the healthcare provider should be able to:

- Describe 3 types of blunt injuries.
- Discuss penetrating injuries.
- Explain the 3 sections of the abdominal cavity and the organs in each.
- Discuss primary and secondary surveys.
- Describe physical signs of bleeding, including Cullen's sign and Grey Turner's sign.
- Describe the 4 classes of hemorrhagic shock.
- Explain the APVU and AMPLE methods of neurological assessment.
Describe key information needed during history for motor vehicle accidents and penetrating trauma.
- Explain FAST and DPL.
- Discuss hepatic trauma, including grading.
- Discuss splenic trauma, including grading.
- Discuss gastric trauma, including grading.
- Discuss intestinal trauma, including at least 6 signs of peritonitis.
- Discuss pancreatic trauma, including grading.
- Discuss bladder trauma and at least 4 indications of bladder rupture.
- Describe the difference between intraperitoneal bladder rupture and extraperitoneal, including implications.
- Discuss renal trauma, including grading.
- Discuss adrenal trauma.
- Discuss aortic trauma.

Introduction



[Wikimedia Commons](#)

Abdominal trauma is responsible for about 10% of all deaths related to trauma. Abdominal trauma may involve penetrating or blunt injuries. Penetrating injuries include gunshot and shrapnel injuries, impalements, and knifings.

- Penetrating injuries often result in injury to hollow organs, such as the intestines. The liver is the most commonly injured solid organ. Gunshot wounds are classified as high energy and may result in extensive damage, especially if the bullet ricochets off of bone. Patients are at increased risk for both hemorrhage and peritonitis, especially with intestinal injury.

The most common injuries from gunshot wounds include:

- Small intestines: 50%.

- Colon: 40%.
- Liver: 30%.
- Vascular structures: 25%.

The most common injuries from stab wounds include:

- Liver: 40%.
 - Small bowel: 30%.
 - Diaphragm: 20%.
 - Colon: 15%.
- Blunt trauma is more common than penetrating and may result from motor vehicle accidents, sports accidents, blows, falls, or explosions. Blunt trauma is more difficult to assess because injuries are less obvious, so massive blood loss may occur before injuries are detected.

There are 3 types of blunt injuries: 1) crush, which results from compression, 2) shear, which involves tearing, and 3) burst, which relates to sudden increased pressure (such as from an explosion). Motor vehicle accidents, the most common cause of blunt abdominal injuries, often result in hepatic injury to the passenger if impact is on the passenger's side and splenic injury to the driver if impact is on the driver's side.

A tremendous force is needed to fracture a pelvis, so any time a trauma patient presents with pelvic trauma, abdominal trauma should be suspected.

Abdominal cavity		
Thoracic abdomen	Liver, spleen, diaphragm, stomach	The ribs provide some protection although the organs are vulnerable to both blunt and penetrating trauma.
Retroperitoneal abdomen	Kidneys, ureters, pancreas, duodenum	Organs are generally better protected than organs in the thoracic abdomen as they are behind other organs.
True abdomen	Small intestines, colon, bladder	The large and small intestines are vulnerable to rupture with blunt trauma. The bladder is somewhat protected behind the symphysis pubis, but a full bladder is especially vulnerable

		to rupture.
--	--	-------------

Common abdominal injuries include:

- Hepatic lacerations.
- Splenic rupture.
- Bladder rupture.
- Mesenteric artery tears.
- Great vessel tears.
- Diaphragmatic rupture.
- Gastric rupture.
- Renal injury.
- Pancreatic injury.
- Retroperitoneal hematoma.

Multiple injuries are common, so any type of abdominal injury should raise suspicions of associated injuries.

Initial assessment

Primary survey: ABCs As with all trauma patients, assessment should begin with the primary survey, during which the patient is at least partially disrobed for examination, placed on monitors (BP, cardiac, pulse oximeter) as indicated, and two large-bore intravenous lines placed. If IV access cannot be achieved, then a large-bore central line may be placed in the femoral vein, subclavian, or internal jugular vein. In some cases, venous cutdown may be indicated. An interosseous line may also be placed, especially in children, if other access is not possible.

Airway, breathing, and circulation (ABCs) are checked immediately, often while resuscitation efforts are occurring. The patient must be assessed for blood loss. With blunt trauma, severe bleeding may occur rapidly, so the patient must be observed for signs of bleeding and blood loss estimated.

Any impaled items, such as a knife stuck into the abdomen, should be stabilized with bulky dressings until scans are completed and/or the patient is taken for surgical removal. Protruding organs or eviscerations should be covered with sterile saline dressings.

Oxygen is usually administered with a non-rebreather mask, and NG tube inserted (if there is no facial trauma), and blood samples (type

and crossmatch and CBC) and urine specimen obtained per indwelling catheter.

Physical signs of internal bleeding include:

- Abdominal pain.
- Guarding, rigidity.
- Bruising, crepitus, swelling (especially across chest and pelvis from seat belt and or shoulder harness).
- Abdominal distention, deformity.
- Tachycardia, hypotension.
- Pallor.
- Evisceration.
- **Cullen's sign:** Bruising about the umbilicus (may indicate hemoperitoneum or retroperitoneal bleeding but may take 12 hours to develop).
- **Grey Turner's sign:** Bruising over flank (may indicate retroperitoneal bleeding but may take 12 hours to develop).
- Hematuria.
- Blood or semen at urethral meatus (from injury to prostate).
- Inability to urinate.

Hemorrhagic shock classification		
Class	Blood loss	Signs and symptoms
I	≤15%	Mild tachycardia (90-100 bpm), localized swelling, and frank bleeding.
II	15 to 25%	Tachycardia, prolonged capillary refill and increased diastolic BP.
III	25 to 50%	Above signs (any) as well as hypotension, confusion, decreased urinary output, and acidosis.
IV	>50%	Hypotension and acidosis unresponsive to resuscitation.

Geriatric patients must be observed carefully as they may have less obvious signs of shock for a variety of reasons. Cardiac response to hypovolemia is often lessened because of myocardial pathology or medications, such as β -blockers and calcium channel blockers.

Metabolic acidosis (decreased serum bicarbonate and increased base deficit (> -6) or increased serum lactate) may result from hemorrhage and hypovolemia.

Secondary survey: DE,

ABPU, and AMPLE The patient is examined more thoroughly to determine which diagnostic tests are indicated (often after initial standard x-rays of later cervical spine, supine chest, and pelvis).

D: Neurological status (D for disability) is assessed. A quick assessment may be done using the **AVPU** method:

- **A:** Is the patient alert?
- **V:** Is the patient verbal?
- **P:** Is the patient responding only to verbal stimuli?
- **U:** Is the patient completely unresponsive.

If the patient is alert (A) and verbal (V), then the AMPLE method may be used to ask a series of basic informational questions:

- **A:** Do you have any allergies?
- **M:** Are you taking any medications?
- **P:** What is your past medical history and (if applicable) are you pregnant?
- **L:** When did you last eat?
- **E:** What events led to the trauma, and when did you have your last tetanus vaccination?

E (for exposure): The patient is completely disrobed, if possible. If there is any danger of cervical trauma or the patient cannot be moved, the clothing should be cut off. If forensics are involved (as in shootings and knifings), care must be used to avoid cutting through or damaging areas of clothing that may provide evidence, such as where a bullet has entered clothing. Protocols for collection of evidence should be carefully followed.

NOTE: In reality, the Primary and secondary surveys may be done concurrently or overlap, depending on the patient's condition.

Key information needed during history	
Motor vehicle accidents	Crash mechanism (what happened). Use of seatbelt and chest restraints. Amount of external damage to the car. Airbag deployment. Integrity of windshield. Deformity of steering wheel Loss of consciousness. Ambulatory status. Other passenger injuries or fatalities. Ejection from vehicle.
Penetrating	Type of weapon involved

trauma	Number of injuries (gunshots, stabbings). Victim's distance from weapon. Patient's position at the time of injury.
---------------	--

Diagnostic procedures: If time is not a critical factor, the **CT scan** is the best tool for assessing all types of abdominal injuries, including bleeding. CT is usually done with contrast. Fast-scanning and image-reconstructing helical CT scanners have reduced the turnaround time for CTs to as little as 10 minutes, but older scanners may require more time.

X-rays, while often done, provide little useful information as they may not identify free air and fluid collections must be large (>800 mL) to be detected by standard x-ray. Angiograms are indicated if injury to vessels is suspected in order to identify the site of bleeding.

However, if the patient is unstable or severe bleeding is suspected, then **focused abdominal sonography in trauma (FAST)** examination should be done.




FAST is able to identify intra-abdominal fluid in about 98% of cases. With multiple trauma patients, FAST may also be used if the CT scan is in use and wait time is extended. However, it's important to remember that while FAST will identify intra-abdominal fluid or bleeding, it is not sufficiently sensitive to show disruption of an internal organ.

Additionally, a single negative finding by FAST does not preclude bleeding or other injury as it is sensitive to >300 mL of blood. For example, a bowel perforation may result in limited intraabdominal fluid in the initial period. Additionally, intracapsular bleeding or delayed organ rupture may not be captured at the time of FAST but may be identified on later examination or CT scan. Additionally, a positive FAST finding does not necessarily indicate a need for surgical intervention.

Diagnostic peritoneal lavage (DPL) may also be used, but FAST is preferred and has generally replaced DPL, as it is relatively fast and

less invasive. DPL is especially insensitive to colonic wounds, which require early diagnosis and treatment.



DPL is done by inserting an abdominal catheter under local anesthetic. Aspiration is done to determine if free blood is present. If aspirate contains <math>< 10\text{ mL}</math> of blood and no evidence of intestinal contents, the aspirate is usually reinjected and 1 L of NS infused with the fluid recollected through gravity drainage.

Positive findings for DPL	
Blunt trauma	RBC/mL: $\geq 100,000$ WBC/mL: ≥ 500 Amylase (IU/L): ≥ 30 Alkaline phosphatase (IU/L): ≥ 3 Intestinal contents: Any amount
Penetrating trauma	RBC/mL (Abdominal stab wound): ≥ 5000 RBC/mL (Gunshot wound): ≥ 5000


Note that a CBC and hemoglobin are not reliable to diagnose blood loss in the initial assessment although serial tests may indicate blood loss or hemodilution related to fluid resuscitation. White blood cell counts with left shift commonly occur after trauma, usually related to demargination.

Usual indications for automatic surgical exploration/repair include:

- Hemodynamic instability.
- Evisceration.
- Peritonitis.
- Free air in abdomen.
- Gunshot wound.

Stab wounds that are superficial may require extending the wound opening and exploring to determine the extent of injury. Laparoscopy may be used to determine if the peritoneum has been violated.

Hepatic trauma



The liver is especially vulnerable to trauma because of its anterior position in the abdomen. It may be lacerated or avulsed by either blunt or penetrating injuries. Liver injury should be suspected if a patient has rib fractures on the right side or has abdominal pain, especially in the right upper quadrant.

After the spleen, hepatic injury is the second most common abdominal injury, with injury to the posterior segment of the right lobe occurring most frequently. Because of this, bleeding may occur into the retroperitoneal area rather than the peritoneal. Injuries that are commonly associated with hepatic trauma include right lung contusion, right pneumothorax, right-sided rib fractures, right kidney and/or

adrenal gland injuries. About 45% of those with hepatic injuries also have injury to the spleen.

Injury of the left lobe is less common and most often associated with direct blow to the epigastric area and is associated with injury to the duodenum, pancreas, and transverse colon.

Mortality rates with hepatic injury are high (8% to 25%), and it is the most common cause of death related to abdominal trauma. With severe injuries, death can occur within minutes. Hepatic injuries can include lacerations, contusions, subcapsular hematoma, and intrahepatic hematoma.

Hepatic injury grading scale	
Grade I	Laceration(s) < 1cm deep. Subcapsular hematoma <1 cm diameter.
Grade II	Laceration(s) 1 to 3 cm deep. Subcapsular or central hematoma 1 to 3 cm diameter.
Grade III	Laceration(s) 3 to 10 cm deep. Subcapsular or central hematoma 3 to 10 cm deep.
Grade IV	Laceration(s) >10 cm deep. Subcapsular or central hematoma >10 cm diameter. Lobar maceration or devascularization.
Grade V	Bilobar tissue maceration or devascularization.


While in the past, surgical management of liver injury was common, it was found that those undergoing surgery tended to have more complications and required more transfusions than those treated more conservatively, so presently only about 20% are treated surgically. In about 70% of cases, bleeding stops spontaneously. Even grade IV injuries may be treated non-surgically if there is no bleeding into the peritoneal or retroperitoneal cavities.

However, patients must be carefully monitored as delayed complications, including hemorrhage, abscess, and biloma (encapsulated collection of bile in the peritoneal cavity), may occur. Additionally, patients may require replacement fluids and blood products, such as plasma and platelets for coagulopathy.

Hepatic contusions usually heal within a week, but subcapsular hematomas may enlarge initially before slowly clearing. Lacerations may require weeks to heal, and small residual bilomas are common.

Splenic trauma

The spleen lies posteriorly in the left upper quadrant, behind the 9th, 10th, and 11th ribs with the convex surface under the left hemidiaphragm. The tail of the pancreas contacts the spleen in about 30% of people and is within 1 cm of the spleen in about 70%. The splenic artery provides the major blood supply to the spleen. The spleen filters approximately 10 to 15% of the total blood volume every minute and usually contains 40 to 50 mL of blood, but the spleen can expand to hold much more blood.



Splenic injuries are the most common solid organ injuries, accounting for about a fourth of total injuries. Splenic injuries may occur along with injuries to other abdominal organs as well. Penetrating injuries are not common, but can occur, especially with gunshot wounds.

Blunt trauma may occur from compression or deceleration injuries, including motor vehicle accidents, blows to the abdomen (kick injuries), and falls.

The most common injury associated with splenic injury is fracture of the left lower rib as this indicates sufficient force to the LUQ to cause organ damage. The triad of symptoms associated with splenic rupture includes:

- Elevation of the left hemidiaphragm.
- Left lower lobe atelectasis.
- Pleural effusion.

Patients may present with various clinical symptoms. Indications of splenic trauma can include left upper quadrant or left flank pain, Kehr sign (referred pain to the left shoulder). Some patients may be

essentially asymptomatic. Indications of shock may be present, including tachycardia, tachypnea, anxiety, and hypotension.

If blood begins to pool in the intraperitoneal area, more diffuse abdominal pain and rebound tenderness may be evident. Hypotension occurring with splenic injury is cause for concern as it may indicate rupture of the spleen with profound hemorrhage.

If shock cannot be compensated, immediate surgical exploration is indicated as the delay needed to confirm hemorrhage by CT scan may result in the patient bleeding out. Patients with compensated shock may be treated with angioembolization if this can be done quickly.

Because of the spleen's importance to the immune system, splenic retention is the goal for those who are hemodynamically stable. Over 90% of children can be treated nonsurgically, regardless of the grade of injury and up to 65% of adults.

Patients who are unstable with suspected injury to the spleen and intra-abdominal hemorrhage may require exploratory laparotomy and repair or removal of the spleen. Additionally, patients with blunt trauma and hemodynamic instability that does not respond to administration of intravenous fluids may be considered to have a life-threatening splenic injury requiring surgery.

Splenic injury grading scale	
Grade I	Laceration(s) < 1cm deep. Subcapsular hematoma <1 cm diameter.
Grade II	Laceration(s) 1 to 3 cm deep. Subcapsular or central hematoma 1 to 3 cm diameter.
Grade III	Laceration(s) 3 to 10 cm deep. Subcapsular or central hematoma 3 to 10 cm deep.
Grade IV	Laceration(s) >10 cm deep. Subcapsular or central hematoma >10 cm diameter.
Grade V	Splenic tissue maceration or devascularization.

The goal with splenic injury is to conserve the spleen whenever possible. Splenectomy increases the risk of postsplenectomy sepsis although it may be necessary with multiple injuries or severe hemodynamic instability.

Postoperatively, recurrent bleeding may occur with splenorrhaphy, especially during the first 24 to 48 hours. Complications in the early

postoperative period for splenectomy or splenorrhaphy can include bleeding, gastric distention, gastric necrosis, pancreatitis, and subphrenic abscess.

Later complications can include deep vein thrombosis and overwhelming post splenectomy infection (OPSI) (usually at 1 to 6 weeks). OPSI can occur within 2 years of splenectomy, especially in children, so prophylactic antibiotics may be given for 2 years. OPSI occurs in about 3% of splenectomy patients with about 50% developing pneumonia or meningitis. Symptoms are often rapid and about half die within 2 days of onset.


Gastric Trauma

Because the stomach has 3 muscle layers, blunt trauma perforations are rare although risk increases if a person suffers a severe blunt force trauma with a full stomach. Forceful blunt trauma may result in rupture of the left hemidiaphragm, causing the stomach to herniate into the left hemithorax. The areas most prone to rupture include:

- Anterior wall: 40%.
- Greater curvature: 23%.
- Lesser curvature: 15%.
- Posterior wall: 15%.

Perforation may result from abdominal impact such as when a pedestrian is struck by a motor vehicle or during a motor vehicle accident in which a person is ejected from the car or the seat belt is improperly applied.

Other injuries can include hematomas and contusions, but these injuries are often essentially asymptomatic and may resolve over time although large hematoma of the distal portion of the stomach may narrow the lumen and prevent emptying. Because the stomach has a rich supply of arteries, damage to the arteries may result in life-threatening hemorrhage.



It's important to note that when blunt trauma is severe enough to cause perforation of the stomach, 95% of patients also have another serious injury. The most common injuries associated with gastric perforation include:

- Spleen: 27% to 43%.
- Left chest: 18% to 29%
- Liver: 18%.
- Small intestine: 18%.

Penetrating trauma of the stomach, such as from a knife or gunshot wound, is more common and should be suspected when penetration is inferior to the nipples or 4th intercostal space anteriorly or inferior to the tips of the scapulae posteriorly.


Symptoms of perforation include severe abdominal pain, abdominal rigidity, hematemesis and bloody nasogastric drainage. While the stomach usually contains few bacteria, with perforation, stomach acids begin to pour into the peritoneal cavity, resulting in chemical peritonitis. Most patients present with shock and pain in the abdomen, but some patients may have no signs of an acute abdomen in the initial period.

Gastric injury grading scale		
I	Intramural hematoma OR superficial laceration	May be observed or drain placed with Lembert, imbricated suturing.
II	<2 cm laceration of pylorus or gastroesophageal junction OR <5 cm laceration of proximal third of stomach OR <10 cm laceration of distal two-thirds of stomach.	Similar to above.
III	>2 cm laceration of pylorus or	Primary surgical repair,

	GE junction OR ≥ 5 cm laceration of proximal third of stomach OR ≥ 10 cm laceration of distal two-thirds of stomach.	pyloroplasty or total gastrectomy (usually for GE junction injury) as indicated.
IV	Perforation or devascularization of <two-thirds of stomach.	Subtotal or total gastrectomy with Roux-en Y or Billroth II reconstruction.
V	Perforation or devascularization of >two-thirds of stomach.	Gastrectomy (Roux-en-Y reconstruction).

Upright x-rays may show free air in the abdomen, but only about 50% to 66% develop enough free air in the abdomen to be detected by upright x-ray. DPL may show food particles or bilious fluids in the abdomen. However, if the patient is hemodynamically stable, CT with contrast provides a definitive diagnosis.

Intestinal trauma



Intestinal trauma may occur as the result of blunt or penetrating trauma, such as from gunshot wounds or knifings. Penetrating trauma may result in evisceration of the small intestines through the abdominal wall. Falls from great heights or crush injuries may result in evisceration through the rectum or perineum. The small intestines are especially vulnerable to penetrating wounds as they cover the

abdominal surface. Motor vehicle accidents also frequently result in injury to the small intestines.

The location of the duodenum and its attachments make it one of the most commonly injured sites, especially with steering wheel injuries that force the duodenum against the spine. Patients often present with back pain when they have duodenal injuries.

Blunt trauma can result from two different types of forces:

- Compression: Increases intraluminal pressure inside of the bowel and compresses the fluid-filled bowel against the vertebrae or other solid structures.
- Deceleration: Stretches and tears the bowel.

Injuries of the small intestine occur about 4 times more frequently than injuries of the colon. Injury to the colon occurs in 2% to 15% of those with blunt abdominal injuries, often motor vehicle accidents. Intestinal trauma may also result from diagnostic or therapeutic procedures, such as colonoscopy, laparoscopy, and radiotherapy, as well as from ingestions, such as from swallowing a toothpick.

Because injury to the colon requires considerable force, other abdominal injuries are frequently present, especially with injury to the transverse colon. Other injuries include:

- Hepatic: 64%.
- Spleen: 52%.
- Small intestinal mesentery: 48%.

Intestinal injuries results in mortality rates of 10% to 30%, but death may be caused by other injuries rather than just the intestinal trauma. Delay in diagnosing and treating intestinal perforation results in mortality rates of 25% to 35%.

As with other abdominal injuries, the CT is the best diagnostic tool and is about 97% accurate in providing evidence of bowel injury; however, CT findings may be compromised if a patient first has DPL, as this procedure may introduce intraperitoneal fluid and air. CT is not always effective in identifying the exact location of an injury to the bowel, and CT in general is less effective in identifying injuries to hollow organs than to solid organs.

Typically, patients have non-specific symptoms or are asymptomatic initially but return with a few hours or a couple of days with signs of peritonitis, including:

- Abdominal distention, rigidity.

- Guarding, rebound tenderness.
- Abdominal pain.
- Lack of bowel sounds, paralytic ileus.
- Elevated WBC counts.
- Fever.
- Tachycardia.
- Dyspnea.
- Nausea and vomiting.

If gross blood is found with a rectal exam, this is usually indicative of severe intestinal injury.

Intestinal perforation may progress to abscess, fistula, and/or sepsis. The site of perforation is an important risk determinant. Little bacteria are present in the proximal segments of the small intestines, but the distal segments contain both aerobic and anaerobic bacteria, which can rapidly result in severe infection. According to some studies, the most commonly perforated areas of the small intestine are the proximal jejunum and the distal ileum.

Treatment of mild intestinal injury may be conservative, but with perforation, surgical intervention with peritoneal lavage is indicated as well as antibiotics. In some cases, primary closure is avoided and the wound left open to heal by secondary intention.


Pancreatic trauma

The pancreas is located in a fairly protected part of the abdomen, high in the retroperitoneum, so when it is injured as the result of blunt trauma, then 90% to 95% of the time, there are injuries to other organs as well. Because of this, injury to the pancreas may be overlooked. Common associated injuries include:

- Liver: 32% to 42%.
- Spleen: 25% to 40%.
- Stomach: 20% to 40%.
- Major vessel: 25% to 35%.
- Thorax: 22% to 31%.
- Intestines: 10% to 29%.
- Central Nervous System: 25%.
- Duodenum: 18%.

While injury to the pancreas occurs in fewer than 10% of all cases of blunt abdominal trauma, the pancreas is injured in 20% to 30% of cases of penetrating trauma. Blunt trauma may result from motor vehicle or pedestrian accidents. Direct blows to the epigastrium may crush the pancreas against the spine.

Penetrating injuries most often are associated with gunshot wounds or stabbings to the back, flank, or abdomen.



Because vascular structures, such as the aorta and the superior mesenteric artery lie close to the head of the pancreas, injury to that area may include adjacent vascular injury and life-threatening hemorrhage.

Symptoms specific to pancreatic trauma are often missing or nonspecific but can include abdominal pain, flank pain, epigastric pain and nausea and vomiting (bile). Diagnosis is often delayed because patients are treated for other injuries and then further testing is done when the patients fail to improve or continue to deteriorate.

As with other injuries, the CT scan provides the most definitive diagnosis as the scan may show lacerations or peripancreatic fluids. Blunt trauma may result in retroperitoneal hematoma and fluid, free abdominal fluid and pancreatic edema. In the early phases after injury, CT scans may not detect injury, so ongoing evaluation is necessary. If the CT scan is inconclusive, then magnetic resonance cholangiopancreatography may be indicated, as it more clearly outlines damage to the ducts.

Standard x-rays are not effective for diagnosis of pancreatic injury but findings of fractures of the lower thoracic or upper lumbar vertebrae should raise suspicion of pancreatic or duodenal injury. FAST may show free abdominal fluid but the position of the pancreas and the overlying of the pancreas by the colonic gas make visualization difficult. DPL has not been found useful because of the retroperitoneal location.

Serum amylase levels have been used to help diagnose pancreatic injury, but studies show that levels don't elevate for about 3 hours, so serum amylase levels prior to 3 hours are not diagnostic. Additionally, patients with brain injuries also have elevations in serum amylase, so if a patient suffers both brain and abdominal trauma, elevation does not necessarily indicate pancreatic trauma. Pancreatic enzymes (amylase and lipase) may be abnormal because of shock rather than direct pancreatic injury, but they may remain normal even with severe pancreatic injury.

There are 3 different grading scales used to classify pancreatic injury. Although they are similar, this can lead to some confusion. The following grading system is that of the Organ Injury Scaling Committee of the American Association for the Surgery of Trauma.

Pancreatic injury grading scale	
I	Simple contusion.
II	Major contusion or laceration without tissue loss or involvement of the main pancreatic duct.
III	Complete transection of the pancreas or a parenchymal injury with involvement of the major duct to the left of the superior mesenteric vein.
IV	Ductal transection or a major parenchymal injury to the right of the superior mesenteric vein.
V	Massive disruption of the head of the pancreas.


Most injuries to the pancreas are relatively minor and can be treated by inserting external drains until healing occurs. Distal pancreatectomy may be necessary with trauma to the body, neck and/or tail with duct disruption. Trauma to the head of the pancreas is usually treated with external drainage even in the presence of ductal damage. Pancreaticoduodenectomy is usually done only when the head of the pancreas has been severely damaged and devitalized.

Post-operative complications can include pancreatic abscess, fistula, pseudocysts, and diabetes mellitus. The most common complication is development of a pancreatic fistula. These usually resolve spontaneously in 1 to 2 weeks with adequate drainage and nutrition. Octreotide is sometimes given perioperatively to prevent fistula development or when a pancreatic fistula occurs. Complications occur in 20% to 40% of those who are treated surgically for pancreatic trauma with 30% of deaths related to sepsis and multi-organ failure. About 10% to 25% develop abscesses. Pseudocysts may develop

weeks or months after trauma. About 18% of those with surgical repair of pancreatic trauma develop mild transient pancreatitis.

Bladder trauma

Urinary Bladder



The urinary bladder is a muscular hollow distensible organ that lies at the floor of the pelvis, protected by the symphysis pubis. The bladder is most vulnerable in children because it is suspended above the symphysis pubis. By about age 20, the bladder descends into the pelvis. Older males are also more vulnerable because an enlarged prostate displaces the bladder upward.

The most common injury is bladder contusion, an incomplete or partial tear of the bladder mucosa, causing localized injury and hematoma. Contusion usually presents as gross hematuria, but the condition usually resolves after a period of bedrest.

Blunt trauma is more likely to result in rupture of the bladder if the bladder is distended, causing it to rise above the symphysis pubis. People who are inebriated at the time of injury in a motor vehicle injury should be carefully examined for possible bladder rupture as excessive drinking often results in a distended bladder. Most blunt trauma results from motor vehicle accidents (85%) with falls (7%) and assaults (6%) causing the rest of the injuries.

Fractures of the pelvis (commonly the anterior pubic arch) or symphysis pubis may also result in rupture either from bone fragments or shear injury. Damage to the bladder may also result from orthopedic trauma when orthopedic pins or screws used to stabilize

pelvic fractures perforate the bladder. Up to a quarter of those with pelvic fractures also have urethral trauma.

The bladder may also be ruptured through penetrating trauma, such as gunshot wounds (85%) and knifings (15%) usually resulting in multiple organ injuries. While non-traumatic bladder perforation can occur, about 82% of bladder rupture relates to external trauma with 60% to 85% from blunt trauma and 15% to 40% from penetrating.

Approximately 50% to 71% of traumatic bladder perforations are extraperitoneal (usually related to pelvic fractures) while 25% to 43% are intraperitoneal (usually related to a direct blow to a distended bladder, such as with a seatbelt injury) and 7% to 14% are combined (often related to gunshot wounds), a much more serious situation with a 60% mortality rate.

Intraperitoneal perforation may remain undiagnosed for extended periods because the urine continues to drain into the abdomen. In this case, the patient may be anuric and may develop electrolyte imbalances as urine is reabsorbed. Urinary ascites may occur with diagnosis per paracentesis.

Associated injuries are common and include:

- Gunshot wounds: $\leq 83\%$ have bowel injuries.
- Stab wounds: 33% have colon injuries.
- Penetrating trauma in general: $\leq 82\%$ have vascular injuries.

While symptoms of bladder injury may be non-specific, a common triad includes hematuria, suprapubic pain or discomfort, and difficulty urinating or anuria. Gross hematuria occurs in about 90% of those with bladder rupture with 88% having pelvic fractures. Other signs of rupture include abdominal distention, guarding, and rebound tenderness. With intraperitoneal rupture, bowel sounds may be absent.


Blood in the urethral meatus may indicate trauma to the urethra, an indication for retrograde urethrography prior to insertion of a Foley catheter because passing a catheter may exacerbate a small tear. If a patient has a posterior urethral injury, a suprapubic catheter should be inserted. Urethral injuries are rare in females but may occur in males because of the longer length and positioning of the urethra. Ureters are rarely injured by blunt trauma but may be damaged by penetrating trauma.

The CT cystogram provides the most definitive diagnosis of bladder rupture. Once a catheter is in place, dilute water soluble contrast is instilled followed by an abdominal/pelvic CT. Note that water soluble contrast is less likely to result in peritonitis if the solution leaks into the abdomen. In some cases, standard cystography may also be indicated.


Treatment depends on the type of rupture:

RUPTURE OF THE BLADDER

Intraperitoneal (A)



Extraperitoneal (B)



- **Extraperitoneal ruptures:** Foley catheter (SP or urethral) for 7 to 10 days with a cystogram to determine if laceration has healed. Almost all heal within 21 days. If extensive extravasation occurs, then surgical exploration may be indicated.


- **Intraperitoneal:** Most require surgical repair as prolonged catheterization alone rarely brings about healing. However, some authorities recommend conservative treatment for small perforations, but there is no consensus regarding this approach.

Note that gunshot wounds are almost always explored surgically, and in that case, even extraperitoneal ruptures may be sutured closed. After surgical exploration, a pelvic drain is usually left in place for 2 to 3 days, and IV antibiotics are administered.

Post-surgical complications can include urinary extravasation (usually treated by extended catheter drainage), wound dehiscence, hemorrhage, infection, and impaired bladder function.

Remember that while only 10% of pelvic fracture patients have a ruptured bladder, 90% of ruptured bladders relate to pelvic fractures, so bladder rupture should always be suspected with pelvic fractures.

Renal trauma



Generally, the kidneys are paired organs in the retroperitoneal space on the posterior abdominal wall extending from the 12th thoracic vertebrae to the 3rd lumbar vertebrae in the adult. The kidneys are well protected by the rib cage and the muscles of the back and abdomen; however, the lower portions of the kidneys extend below

the 12th ribs. Fatty deposits that protect them against jarring surround the kidneys. An adrenal gland sits on top of each kidney.

The kidneys receive 20% to 25% of the total cardiac output, and all of the body's blood circulates through the kidneys about 12 times each hour, so damage to the kidney's vasculature may result in hemorrhage.

About 10% of those with abdominal trauma sustain renal injuries with blunt trauma injuries about 9 times more frequent than penetrating trauma injuries. Blunt injuries include renal contusion, renal laceration, and renal vascular injury. Blunt injuries usually result from motor vehicle accidents, falls, and pedestrian accidents or sport injury that result in a direct blow to the flank area.

Symptoms may be very nonspecific but can include abdominal or flank pain and gross or microscopic blood in the urine. In the renal injury grading system, grades I and II are classified as minor injuries and grades III, IV, and V as major. Major renal trauma is usually caused by penetrating injuries (40%) rather than blunt (15%).

Renal injury grading system	
I	Contusion: Microscopic or gross hematuria. Urological studies normal. Hematoma: Subcapsular, nonexpanding without parenchymal laceration.
II	Hematoma: Nonexpanding perirenal hematoma confined to renal retroperitoneum. Laceration: <1cm parenchymal depth of renal cortex without urinary extravasation.
III	Laceration: >1 cm depth of renal cortex, without collecting system rupture or urinary extravasation.
IV	Laceration: Parenchymal laceration extending through the renal cortex, medulla, and collecting system. Vascular: Main renal artery or vein injury with contained hemorrhage.
V	Laceration: Completely shattered kidney. Vascular: Avulsion of renal hilum, which devascularizes kidneys.

Diagnosis is based on a number of factors. Urinalysis is standard, but some types of renal injury (avulsion, renal artery laceration) may not result in hematuria, so the absence of blood does not rule out damage to the kidneys. In the past the intravenous pyelogram (IVP) was used


for diagnosis, but the CT scan with contrast is now the procedure of choice.

While in the past surgical repair was the standard, with low-grade blunt trauma, the kidneys actually usually heal with bedrest and observation. Most injuries to the kidneys (grades I to IV) are managed non-surgically, but patients must be monitored carefully after treatment as late bleeding may occur, and some patients develop hypertension because of kidney damage.

With high-grade trauma or penetrating injuries, surgical exploration may be indicated, especially if the patient has other abdominal injuries and is hemodynamically unstable. Only about 9% of renal injuries require surgical intervention and about 11% of these result in nephrectomy, usually because of hemorrhage or severe renovascular injury. Complications can also include perirenal abscess and or urinary leakage.

Adrenal trauma

Adrenal Gland



Trauma to the adrenal glands is rare (<1% to 2%) because they are well protected under the rib cage, but injuries do occur, especially with multiple organ traumas. Adrenal gland trauma is associated with mortality rates about 5 times higher than if there is no adrenal gland trauma. Injury may result from blunt trauma (most commonly motor vehicle accidents), or penetrating trauma.

Adrenal trauma is associated with:

- Hepatic injury.
- Splenic injury.
- Renal injury.
- Rib fractures.
- Lumbar fractures.
- Pelvic fractures.

In many cases, patients also have multiple other injuries, such as head injuries and injuries to the extremities.

Hematoma or laceration of the adrenal gland often is found incidentally with CT scan to evaluate other abdominal injuries. One study found that left adrenal hematomas were most commonly associated with left rib fractures and splenic and left renal injuries while right adrenal hematomas were associated with right rib fracture and hepatic and right renal injuries.

One study of pediatric patients with adrenal trauma showed that those with unilateral injury rarely showed evidence of adrenal insufficiency, but adrenal insufficiency should be suspected in bilateral injuries.

About 78% of adrenal hematomas occur on the right, possibly because the area about the gland is more confined because of the mass of the liver on the right. Additionally, high deceleration pressures may be transmitted through the inferior vena cava and the short right adrenal gland. The right adrenal gland lies directly behind the inferior vena cava.


Diagnosis is by CT but may be impacted by the presence of benign adenomas, which are common but can be misdiagnosed as hematomas. FAST may also be used as a screening tool.

Treatment is usually conservative as hematomas usually resolve over time. In severe injuries, such as with gunshot wounds, adrenalectomy may be indicated. Hemorrhage may require transfusions. Acute adrenal insufficiency (usually related to bilateral injury and hemorrhage) must be quickly diagnosed and treated with glucocorticoids.

Aortic trauma


The aorta can be injured with both blunt and penetrating trauma. Tearing may occur from sudden deceleration, as in motor vehicle accidents, and impact with the steering wheel may result in

compression that ruptures the aorta. About 80% to 90% of great vessel trauma is fatal with 15% of deaths related to motor vehicle accidents caused by aortic trauma. Those with small tears or partial thickness tears may survive.



WORLDVET.V.ORG®

A free rupture typically present with widened mediastinum,



hemothorax and hemodynamic instability and is almost always fatal; however, a controlled rupture in which there is no hemothorax and the patient remains hemodynamically stable has about a 90% survival rate with prompt surgical intervention.

The most common site of aortic trauma is distal to the

left subclavian at the level of the ligamentum arteriosum as this is the point of maximum stress and tearing from deceleration forces. Therefore, the aorta is at risk in frontal and side impacts as well as falls from heights.

If aortic injury is not treated promptly the patient may develop hypoxia and hypovolemia with anoxic encephalopathy and ischemic damage to other viscera.

Diagnosis of aortic injury is through CT scan. If the scan is not definitive, than an aortogram may be done to identify small tears. In some cases intravascular ultrasound (IVUS) or transesophageal echocardiography (TEE) may be done instead of aortogram, as these procedures are less invasive.

CT scan may be preceded by chest x-ray (with head elevated) and insertion of an NG tube and/or chest tube. The CT may show a contained rupture, which is referred to as a traumatic pseudoaneurysm.



Chest x-ray is not diagnostic but may show an abnormal mediastinum, distortion of the aorta, and depressed left main stem bronchus as well as deviation of a nasogastric tube. These findings are suggestive of aortic injury. Bleeding may result from tears of mediastinal veins, but this is an indirect indication of possible aortic injury.

If patients with suspected aortic injury are hemodynamically unstable, priority is given to controlling hemorrhage while avoiding over-resuscitation. FAST or DPL may be utilized to identify concealed areas of hemorrhage. Patients with aortic tear and impending rupture may develop a cyclical pattern of responding to fluid resuscitation and then

exhibiting hypotension. A cycle of repeated fluid resuscitations can lead to rupture, especially if other signs, such as widened mediastinum and left-sided hemothorax, are present.

Aortic injury is associated with diaphragmatic rupture, so any patient with aortic injury should be assessed for injury to the diaphragm and *vice versa*.

Aortic injuries are almost always repaired surgically. In some cases thoracic endovascular aortic repair (TEVAR) with insertion of stents may be used. Recent studies indicate that delaying surgery for up to 4 days in non-acute cases, allowing other injuries to be treated and the patient stabilized, has resulted in a lower overall mortality rate than rushing patients into surgery.

If the patient must undergo a craniotomy or exploratory laparotomy because of other injuries and the aortic injury is not acute, β -blockers may be administered to reduce heart rate and force of contractions. Nitroprusside may also be used to control blood pressure and prevent rupture.

Conclusion

Time is a critical factor with abdominal trauma. Patients must be assessed and evaluated immediately, and a finding of pelvic fracture or trauma to one abdominal organ should always raise suspicion of associated injuries.

The diaphragm is the broad muscle that separates the thoracic cavity from the abdominal cavity. The diaphragm is rarely (<5%) injured with blunt trauma, and most of these injuries result from motor vehicle accidents with lateral impact 3 times more likely to cause tears than frontal impacts because of distortion of the chest wall and shearing. Diaphragm injuries may occur from either gunshot wounds or stabbings.

Diaphragmatic injuries rarely occur in isolation so they may be overlooked when attention focuses on associated abdominal injuries, but diaphragmatic injury should be suspected in those with abdominal trauma presenting with ventilatory compromise.

References

- Abdominal trauma: Role of CT. (n.d.). *The Radiology Assistant*. Retrieved November 9, 2011, from <http://www.radiologyassistant.nl/en/466181ff61073>
- American Academy of Orthopaedic Surgeons. (2009). *Advanced Assessment and Treatment of Trauma*. Sudbury, MA: AAOS.
- Bjerke, HS. (2009, August 6). Splenic rupture. *Medscape Reference*. Retrieved November 9, 2011, from <http://emedicine.medscape.com/article/432823-overview>
- Bjerke, HS. (2010, January 2). Pancreatic Trauma. *Medscape Reference*. Retrieved November 9, 2011, from <http://emedicine.medscape.com/article/433177-overview>
- Chest trauma: Traumatic aortic injury. (2004, April). *Trauma.org*. Retrieved November 9, 2011, from <http://www.trauma.org/archive/thoracic/CHESTaorta.html>
- Degiannis, E, Glapa, M, Loukogeorgakis, SP, & Smith, MD. (2008). Management of pancreatic trauma. *International Journal of the care of the Injured*, 39: 21-29. Retrieved November 9, 2011, from <http://www.smmemx.org/Documentos/Trauma/PancreaticTrauma.pdf>
- Fotheringham, T. (2008). Pancreatic injuries. *Nordic Trauma Society*. Retrieved November 9, 2011, from <http://www.nordictraumarad.com/Syllabus08/pancreas.pdf>
- Havens, J, & Lieberman, G. (2002, November 17). Blunt thoracic aortic trauma. *Harvard Medical School*. Retrieved November 9, 2011, from <http://eradiology.bidmc.harvard.edu/LearningLab/cardio/Havens.pdf>
- Hepatic trauma: Liver laceration. (2004). *LearningRadiology.com*. Retrieved November 9, 2011, from <http://www.learningradiology.com/notes/ginotes/livertraumapage.htm>
- Kawashima, A, et al. (2001, May). Imaging of renal trauma: A comprehensive review. *RadioGraphics*. Retrieved November 9, 2011, from <http://radiographics.rsna.org/content/21/3/557.full>
- Kidney (renal) trauma. (2011, January). *AUA Foundation*. Retrieved November 9, 2011, from <http://www.urologyhealth.org/urology/index.cfm?article=61>
- Lusaya, DG. (2011, October 20). Renal trauma. *Medscape Reference*. Retrieved November 9, 2011, from <http://emedicine.medscape.com/article/440811-overview#showall>
- McPhee, SM, & Papadakis, MA. (2009). *Current Medical Diagnosis & Treatment*. San Francisco: McGraw Hill Medical.

- Mick, MW, Peter, JR, Egan, D, & Nadel, ES. (2006). *Blueprints Emergency Medicine*. 2nd ed. Baltimore, MD: Lippincott Williams & Wilkins.
- Mitchell, EL, & Medson, R. (2005). *Introduction to Emergency Medicine*. Philadelphia: Lippincott Williams & Wilkins.
- Mukhopadhyay, M. (2009). Intestinal injury from blunt abdominal trauma: A study of 47 cases. *OMJ*, 24: 256-459. Retrieved November 9, 2011, from http://www.omjournal.org/OriginalArticles/FullText/200910/FT_IntestinalInjuryfromBlunAbdominalTraumaAStudy.html
- Rackley, R. (2009, August 17). Bladder trauma. *Medscape Reference*. Retrieved November 9, 2011, from <http://emedicine.medscape.com/article/441124-overview>
- Rana, AI, et al. (2004, March). Adrenal gland hematomas in trauma patients. *Radiology*, 230: 660-675. Retrieved November 9, 2011, from <http://radiology.rsna.org/content/230/3/669.full>
- Seldman, C. (n.d.). Renal trauma. *Trauma.org*. Retrieved November 9, 2011, from <http://www.trauma.org/archive/abdo/renal/intro.html>
- Smeltzer, SC, Bare, BG, Hinkle, JL, & Cheever, KH. (2008). *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*, 11 ed., Philadelphia: Wolters Kluwer/Lippincott, Williams, & Wilkins.
- Stanton-Maxey, KJ. (2011, August 9). Penetrating abdominal trauma. *Medscape Reference*. Retrieved November 9, 2011, from <http://emedicine.medscape.com/article/2036859-overview#a0104>
- Tritos, NA. (2011, July 13). Adrenal hemorrhage. *Medscape Reference*. Retrieved November 9, 2011, from <http://emedicine.medscape.com/article/126806-overview>